


Application No. 09/843,161
 Amendment dated October 14, 2005
 Page 2

Amendments to the Claims:

The following claims will replace all prior versions of the claims in this application (in the unlikely event that no claims follow herein, the previously pending claims will remain):

1-60. (Cancelled).


61. 1. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of a compound represented by the following formula 1:



[1]

in a physiologically acceptable carrier;

wherein Z is



n, m, q and r independently represent Integers from zero to 4 provided that n + m < 4 and q + r < 4; p and s independently represent integers from zero to 5 provided that p + s < 5; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

Application No. 09/843,16.
Amendment dated October 14, 2005
Page 3

R and R' each independently represent a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; -CONR₂''' halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR''' halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; or linear or branched C₂-C₂₀ alkenyl;

R'''' independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; or optionally substituted C₁-C₂₀ alkoxy;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy carbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; or aralkanoyl;

or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-.

62.2. (Currently Amended) A method according to claim 61, wherein R'
represents -CO₂R'''', CO₂Z' or -CONR₂'''.

63-64. (Cancelled).

Application No. 09/843,167
Amendment dated October 14, 2005
Page 4

66. 3. (Currently Amended) A method according to claim 622, wherein X is -S- and X' is >NH.

71. 4. (Currently Amended) A method according to claim 622, wherein at least two A groups represent a hydrogen atom.

115. 5. (Currently Amended) A method according to claim 622 wherein R' represents -CO₂R''.

67. 6. (Currently Amended) A method according to claim 4155, wherein X is -S- and X' is >NH.

125. 7. (Currently Amended) A method of claim 676 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

116. 9. (Currently Amended) A method according to claim 4155 wherein R'' represents methyl.

132. 8. (Currently Amended) A method according to claim 4257 wherein A', A'', B' and B'' all represent hydrogen atoms.

73. 10. (Currently Amended) A method according to claim 1169 wherein said A group represents methoxy.

117. 11. (Currently Amended) A method according to claim 622 wherein R' represents -CO₂Z'.

68. 12. (Currently Amended) A method according to claim 41711, wherein X is -S- and X' is >NH.

Application No. 09/843,161
Amendment dated October 14, 2005
Page 5

426. 13. (Currently Amended) A method of claim 6812 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

418. 14. (Currently Amended) A method according to claim 11711 wherein Z' is a pharmaceutically acceptable counter ion.

74. 15. (Currently Amended) The method of claim 11814 wherein said pharmaceutically acceptable counter ion is selected from sodium, potassium, calcium, magnesium, ammonium, tromethamine, or tetramethylammonium.

419. 16. (Currently Amended) A method according to claim 622 wherein R' represents -CONR₂"".

420. 17. (Currently Amended) A method according to claim 11916 wherein at least one R"" independently represents a hydrogen atom, methyl or methoxy.

421. 18. (Currently Amended) A method according to claim 11916, wherein both R"" are the same and represent a hydrogen atom, methyl, or methoxy.

422. 19. (Currently Amended) A method according to claim 11916, wherein X is -S- and X' is >NH.

427. 20. (Currently Amended) A method of claim 11916 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

428. 21. (Currently Amended) A method of claim 622 wherein at least two A groups represent methoxy.

470. 22. (Currently Amended) A method of claim 622 wherein said compound is selected from the group consisting of 3-(3,5-dimethoxyphenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-acrylic acid,

Application No. 09/843,16/
Amendment dated October 14, 2005
Page 6

3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-acrylamide,
3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-N,N-dimethyl-acrylamide,
3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-N-methoxy,-N-methyl-acrylamide,
3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylidenemethyl)-phenoxy]-phenyl}-propionic acid methyl ester,
3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylidenemethyl)-phenoxy]-phenyl}-acrylic acid methyl ester,
3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-propionic acid,
3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylidenemethyl)-phenoxy]-phenyl}-propionic acid,
3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylidenemethyl)-phenoxy]-phenyl}-acrylic acid, and
3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-propionic acid methyl ester.

65. 23. (Currently Amended) A method according to claim 611, wherein X is -S- and X' is >NH.

69. 24. (Currently Amended) A method according to claim 611, wherein the bond labeled "a" in formula I represents a single bond.

124. 25. (Currently Amended) A method according to claim 6924 wherein the bond labeled "b" in formula I represents a double bond.

70. 26. (Currently Amended) A method according to claim 611, wherein at least one A group represents methoxy.

72. 27. (Currently Amended) A method according to claim 7026, wherein at least two A groups represent a hydrogen atom.

Application No. 09/843,161,
Amendment dated October 14, 2005
Page 7

75. 28. (Currently Amended) The method of claim 7026 wherein said pharmaceutically acceptable counter ion is selected from sodium, potassium, calcium, magnesium, ammonium, tromethamine, or tetramethylammonium.


123. 29. (Currently Amended) A method according to claim 641 wherein the bond labeled "b" in formula I represents a double bond.

129. 30. (Currently Amended) A method of claim 641 wherein A' and B' represent hydrogen atoms.

130. 31. (Currently Amended) A method of claim 641 wherein A" and B" represent hydrogen atoms.

131. 32. (Currently Amended) A method of claim 641 wherein A', A", B' and B" all represent hydrogen atoms.

76. 33. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of a compound represented by the following formula 1:




[1]

in a physiologically acceptable carrier;

wherein Z is

Application No. 09/843,167
 Amendment dated October 14, 2005
 Page 8



n, m, q and r independently represent integers from zero to 4 provided that n + m < 4 and q + r < 4; p and s independently represent integers from zero to 5 provided that p + s < 5; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R and R' each independently represent a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; -CONR₂''''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; or linear or branched C₂-C₂₀ alkenyl;

R'''' independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; or optionally substituted C₁-C₂₀ alkoxy;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

Application No. 09/843,167
Amendment dated October 14, 2005
Page 9

A, and A' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy carbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

A" independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy carbonyl; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; or halo;

B, B' and B" each independently represent; C₂-C₂₀ alkenoyl; aroyl; or aralkanoyl;

or A and B jointly, A' and B' jointly, or A" and B" jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-.

77. 34. (Currently Amended) A method according to claim 7633, wherein R' represents -CO₂R'', -CO₂Z' or -CONR₂'''.

84. 35. (Currently Amended) A method according to claim 7734, wherein X is -S- and X' is >NH.

85. 36. (Currently Amended) A method according to claim 7734, wherein at least one A group represents methoxy.

87. 37. (Currently Amended) A method according to claim 8536, wherein at least two A groups represent a hydrogen atom.

90. 38. (Currently Amended) The method of claim 8536 wherein said pharmaceutically acceptable counter ion is selected from sodium, potassium, calcium, magnesium, ammonium, tromethamine, or tetramethylammonium.

86. 39. (Currently Amended) A method according to claim 7734, wherein at least two A groups represent a hydrogen atom.

Application No. 09/843,167
Amendment dated October 14, 2005
Page 10

133. 40. (Currently Amended) A method according to claim 7734 wherein R' represents -CO₂R''.

78. 41. (Currently Amended) A method according to claim 43340 wherein R'' represents methyl.

82. 42. (Currently Amended) A method according to claim 43340, wherein X is -S- and X' is >NH.

134. 43. (Currently Amended) A method according to claim 43340 wherein R'' represents methyl.

88. 44. (Currently Amended) A method according to claim 43443 wherein said A group represents methoxy.

142. 45. (Currently Amended) A method of claim 43340 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

149. 46. (Currently Amended) A method according to claim 43340 wherein A', A'', B' and B'' all represent hydrogen atoms.

135. 47. (Currently Amended) A method according to claim 7734 wherein R' represents -CO₂Z'.

83. 48. (Currently Amended) A method according to claim 43547, wherein X is -S- and X' is >NH.

143. 49. (Currently Amended) A method of claim 43547 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

436. 50. (Currently Amended) A method according to claim 43547 wherein Z' is a pharmaceutically acceptable counter ion.

Application No. 09/843,161,
Amendment dated October 14, 2005
Page 11

89. 51. (Currently Amended) The method of claim 43650 wherein said pharmaceutically acceptable counter ion is selected from sodium, potassium, calcium, magnesium, ammonium, tromethamine, or tetramethylammonium.

437. 52. (Currently Amended) A method according to claim 43650 wherein R' represents -CONR₂"".

79. 53. (Currently Amended) A method according to claim 43752 wherein both R'" are the same and represent a hydrogen atom, methyl, or methoxy.

138. 54. (Currently Amended) A method according to claim 43752 wherein at least one R'" independently represents a hydrogen atom, methyl or methoxy.

139. 55. (Currently Amended) A method according to claim 43752, wherein both R'" are the same and represent a hydrogen atom, methyl, or methoxy.

444. 56. (Currently Amended) A method of claim 43752 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

445. 57. (Currently Amended) A method of claim 7734 wherein at least two A groups represent methoxy.

80. 58. (Currently Amended) A method according to claim 7633, wherein X is -S- and X' is >NH.

84. 59. (Currently Amended) A method according to claim 7633, wherein the bond labeled "a" in formula I represents a single bond.

444. 60. (Currently Amended) A method according to claim 8459 wherein the bond labeled "b" in formula I represents a double bond.

Application No. 09/843,167
 Amendment dated October 14, 2005
 Page 12


140. 61. (Currently Amended) A method according to claim 7633 wherein the bond labeled "b" in formula I represents a double bond.

146. 62. (Currently Amended) A method of claim 7633 wherein A' and B' represent hydrogen atoms.

147. 63. (Currently Amended) A method of claim 7633 wherein A" and B" represent hydrogen atoms.

148. 64. (Currently Amended) A method of claim 7633 wherein A', A", B' and B" all represent hydrogen atoms.


91. 65. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of a compound represented by the following formula 1:



[1]


in a physiologically acceptable carrier;

wherein Z is



or

Application No. 09/843,167
 Amendment dated October 14, 2005
 Page 13



n, m, q and r independently represent integers from zero to 4 provided that n + m < 4 and q + r < 4; p and s independently represent integers from zero to 5 provided that p + s < 5; a, b and c represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R"'; -NH₂; -NHR"'; -NR₂"; -OH; -OR"'; -CONR₂"; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R"'; -NH₂; -NHR"'; -NR₂"; -OR"'; -CONR₂"; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R" independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R"'; -NH₂; -NHR"'; -NR₂"; -OH; -OR"'; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; or linear or branched C₂-C₂₀ alkenyl;

R"" independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; or optionally substituted C₁-C₂₀ alkoxy;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

Application No. 09/843,161
Amendment dated October 14, 2005
Page 14

A, A' and A" each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy carbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B" each independently represent; C₂-C₂₀ alkenoyl; aroyl; or aralkanoyl;

or A and B jointly, A' and B' jointly, or A" and B" jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-.

92. 66. (Currently Amended) A method according to claim 9165, wherein R' represents -CO₂R'', CO₂Z' or -CONR₂'''.

96. 67. (Currently Amended) A method according to claim 9266, wherein X is -S- and X' is >NH.

99. 68. (Currently Amended) A method according to claim 9266, wherein the bond labeled "a" represents a single bond.

159. 69. (Currently Amended) A method according to claim 9968 wherein the bond labeled "b" in formula I represents a double bond.

100. 70. (Currently Amended) A method according to claim 9266, wherein at least one A group represents methoxy.

102. 71. (Currently Amended) A method according to claim 10070, wherein at least two A groups represent a hydrogen atom.

105. 72. (Currently Amended) The method of claim 10070 wherein said pharmaceutically acceptable counter ion is selected from sodium, potassium, calcium, magnesium, ammonium, tromethamine, or tetramethylammonium.

Application No. 09/843,167
Amendment dated October 14, 2005
Page 15

101. 73. (Currently Amended) A method according to claim 9266, wherein at least two A groups represent a hydrogen atom.

150. 74. (Currently Amended) A method according to claim 9266 wherein R' represents -CO₂R'''.

93. 75. (Currently Amended) A method according to claim 15074 wherein R''' represents methyl.

97. 76. (Currently Amended) A method according to claim 15074, wherein X is -S- and X' is >NH.

151. 77. (Currently Amended) A method according to claim 15074 wherein R''' represents methyl.

103. 78. (Currently Amended) A method according to claim 15177 wherein said A group represents methoxy.

160. 79. (Currently Amended) A method of claim 15074 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

152. 80. (Currently Amended) A method according to claim 9266 wherein R' represents -CO₂Z'.

98. 81. (Currently Amended) A method according to claim 15280, wherein X is -S- and X' is >NH.

153. 82. (Currently Amended) A method according to claim 15280 wherein Z' is a pharmaceutically acceptable counter ion.

104. 83. (Currently Amended) The method of claim 15382 wherein said pharmaceutically acceptable counter ion is selected from sodium, potassium, calcium, magnesium, ammonium, tromethamine, or tetramethylammonium.

Application No. 09/843,161
Amendment dated October 14, 2005
Page 16

161. 84. (Currently Amended) A method of claim 15280 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

154. 85. (Currently Amended) A method according to claim 9266 wherein R' represents -CONR₂"".

94. 86. (Currently Amended) A method according to claim 15485 wherein both R"" are the same and represent a hydrogen atom, methyl, or methoxy.

155. 87. (Currently Amended) A method according to claim 15485 wherein at least one R"" independently represents a hydrogen atom, methyl or methoxy.

156. 88. (Currently Amended) A method according to claim 15587 wherein both R"" are the same and represent a hydrogen atom, methyl, or methoxy.

157. 89. (Currently Amended) A method according to claim 15485, wherein X is -S- and X' is >NH.


162. 90. (Currently Amended) A method of claim 15485 wherein the bond labeled "b" in formula I represents a double bond and the bond labeled "a" in formula I represents a single bond.

95. 91. (Currently Amended) A method according to claim 9165, wherein X is -S- and X' is >NH.

158. 92. (Currently Amended) A method according to claim 9165 wherein the bond labeled "b" in formula I represents a double bond.

166. 93. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of a compound represented by the following formula 1:


Application No. 09/843,161
 Amendment dated October 14, 2005
 Page 17



[1]

in a physiologically acceptable carrier;

wherein Z is



n, m, q and r independently represent integers from zero to 4 provided that n + m < 4 and q + r < 4; p and s independently represent integers from zero to 5 provided that p + s < 5; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R and R' each independently represent a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

Application No. 09/843,161
Amendment dated October 14, 2005
Page 18

R''' independently represents a linear or branched C₁-C₂₀ alkyl; or linear or branched C₂-C₂₀ alkenyl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, A' and A" each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy carbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B" each independently represent; C₂-C₂₀ alkenoyl; aroyl; or aralkanoyl;

or A and B jointly, A' and B' jointly, or A" and B" jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-.

407. 94. (Currently Amended) A method according to claim 40693, wherein R' represents -CO₂R''' or CO₂Z'.

408. 95. (Currently Amended) A method according to claim 40794, wherein X is -S- and X' is >NH.

409. 96. (Currently Amended) A method according to claim 40794 wherein R' represents -CO₂R'''.

410. 97. (Currently Amended) A method according to claim 40396 wherein R''' represents methyl.

411. 98. (Currently Amended) A method according to claim 40396, wherein X is -S- and X' is >NH.

412. 99. (Currently Amended) A method according to claim 40794 wherein R' represents -CO₂Z'.


Application No. 09/843,161
 Amendment dated October 14, 2005
 Page 19

166. 100. (Currently Amended) A method according to claim 16599 wherein Z' is a pharmaceutically acceptable counter ion.

168. 101. (Currently Amended) A method according to claim 16599, wherein X is -S- and X' is >NH.

108. 102. (Currently Amended) A method according to claim 10693, wherein X is -S- and X' is >NH.


110. 103. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of a compound represented by the following formula 1:



[1]

in a physiologically acceptable carrier;

wherein Z is



n, m, q and r independently represent integers from zero to 4 provided that n + m < 4 and q + r < 4; p and s independently represent integers from zero to 5 provided that p + s < 5; a and b represent double bonds which may be present or absent; when

DCDB01 20767684.1 14-Oct-05 12:12

Application No. 09/843,161,
Amendment dated October 14, 2005
Page 20

present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R and R' each independently represent a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R"'; -NH₂; -NHR"'; -NR₂"; -OH; -OR"'; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R" independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R"'; -NH₂; -NHR"'; -NR₂"; -OH; -OR"'; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; or linear or branched C₂-C₂₀ alkenyl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, and A' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy carbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

A" independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy carbonyl; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; or halo;

B, B' and B" each independently represent; C₂-C₂₀ alkenoyl; aroyl; or aralkanoyl;

or A and B jointly, A' and B' jointly, or A" and B" jointly, independently represent a methylenedioxy or ethylenedioxy group; and


X and X' independently represent >NH, >NR'', -O-, or -S-.

114. 104. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a

Application No. 09/843,161
 Amendment dated October 14, 2005
 Page 21

therapeutically effective amount of a compound represented by the following formula


1:




[1]

in a physiologically acceptable carrier;

wherein Z is



or



n, m, q and r independently represent integers from zero to 4 provided that n + m < 4 and q + r < 4; p and s independently represent integers from zero to 5 provided that p + s < 5; a, b and c represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR''';

DCDB01 20767684.1 14-Oct-05 12:12

Application No. 09/843,161,
Amendment dated October 14, 2005
Page 22

halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R"'; -NH₂; -NHR"'; -NR₂"; -OR"'; -CONR₂"; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R" Independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R"'; -NH₂; -NHR"'; -NR₂"; -OH; -OR"'; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; or linear or branched C₂-C₂₀ alkenyl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, A' and A" each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy carbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B" each independently represent; C₂-C₂₀ alkenoyl; aroyl; or aralkanoyl;

or A and B jointly, A' and B' jointly, or A" and B" jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-.

112. 105. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of 3-(3,5-dimethoxyphenyl)-2-{4-[2,4-dioxothiazolidin-5-ylmethyl]-phenoxy}-phenyl}-acrylic acid in a physiologically acceptable carrier.

Application No. 09/843,161.
Amendment dated October 14, 2005
Page 23

413. 106. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of 3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-acrylamide in a physiologically acceptable carrier.

414. 107. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of 5-(4-(4-(1-carbomethoxy-2-(3,5-dimethoxy phenyl)-ethenyl)-phenoxy)-benzyl)-2,4-thiazolidinedione in a physiologically acceptable carrier.

469. 108. (Currently Amended) A method of treating diabetes comprising the steps of administering to a subject suffering from a diabetic condition, a therapeutically effective amount of 3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-N,N-dimethyl-acrylamide, a physiologically acceptable carrier.